

Fourth ABC Index and Fifth GA Index of Certain Special Molecular Graphs

Yun Gao, Li Liang, Wei Gao

Abstract— Several chemical indices have been introduced in theoretical chemistry to measure the properties of molecular structures, such as atom bond connectivity index and geometric-arithmetic index. In this paper, we present the fourth atom bond connectivity index and fifth geometric-arithmetic index of fan molecular graph, wheel molecular graph, gear fan molecular graph, gear wheel molecular graph, and their r -corona molecular graphs.

Index Terms— fourth atom bond connectivity index, fifth geometric-arithmetic index, r -corona molecular graph

I. INTRODUCTION

Atom bond connectivity index, geometric-arithmetic index and other chemical indices are introduced to reflect certain structural features of organic molecules (See Yan et al., [1-2], Gao et al., [3-4], Gao and Shi [5], Gao and Wang [6], Xi and Gao [7-8], Xi et al., [9], Gao et al., [10] for more detail). The notation and terminology used but undefined in this paper can be found in [11].

M. Ghorbani et al., introduced the fourth Atom-Bond Connectivity index $ABC_4(G)$ [12] as

$$ABC_4(G) = \sum_{uv \in E(G)} \sqrt{\frac{S(u) + S(v) - 2}{S(u)S(v)}},$$

where $S(v) = \sum_{uv \in E(G)} d(u)$. More results on fourth ABC index can refer to [13-15].

Similarity, the fifth geometric-arithmetic index was denoted by [16] as

$$GA_5(G) = \sum_{uv \in E(G)} \frac{2\sqrt{S(u)S(v)}}{S(u) + S(v)}.$$

More conclusions on GA_5 can refer to [17-18].

We define the general fifth geometric-arithmetic index as $GA_5^k(G) = \sum_{uv \in E(G)} \left(\frac{2\sqrt{S(u)S(v)}}{S(u) + S(v)} \right)^k$.

Let P_n and C_n be path and cycle with n vertices. The molecular graph $F_n = \{v\} \vee P_n$ is called a fan molecular graph and the molecular graph $W_n = \{v\} \vee C_n$ is called a wheel molecular graph. Molecular graph $I_r(G)$ is called r -crown molecular graph of G which splicing r hang edges for every vertex in G . By adding one vertex in every two adjacent

vertices of the fan path P_n of fan molecular graph F_n , the resulting molecular graph is a subdivision molecular graph called gear fan molecular graph, denote as \tilde{F}_n . By adding one vertex in every two adjacent vertices of the wheel cycle C_n of wheel molecular graph W_n , The resulting molecular graph is a subdivision molecular graph, called gear wheel molecular graph, denoted as \tilde{W}_n .

In this paper, we present the fourth Atom-Bond Connectivity index of $I_r(F_n)$, $I_r(W_n)$, $I_r(\tilde{F}_n)$ and $I_r(\tilde{W}_n)$. Also, the fifth geometric-arithmetic index and its general version of $I_r(F_n)$, $I_r(W_n)$, $I_r(\tilde{F}_n)$ and $I_r(\tilde{W}_n)$ are derived.

II FOURTH ATOM-BOND CONNECTIVITY INDEX

Theorem1.

$$\begin{aligned} ABC_4(I_r(F_n)) &= \\ r \sqrt{\frac{(4n-4)+r(n+2)}{(n+r)((3n-2)+r(n+1))}} &+ \\ 2 \sqrt{\frac{(4n-1)+r(n+4)}{((3n-2)+r(n+1))(n+3r+3)}} & \\ + 2 \sqrt{\frac{(4n+1)+r(n+5)}{((3n-2)+r(n+1))(n+4r+5)}} + (n-4) \sqrt{\frac{(4n+2)+r(n+5)}{((3n-2)+r(n+1))(n+4r+6)}} & \\ + 2 \sqrt{\frac{2n+8r+9}{(n+4r+5)(n+4r+6)}} + \frac{n-3}{n+4r+6} \sqrt{2n+8r+10} &+ \\ 2r \sqrt{\frac{n+4r+3}{(n+3r+3)(2+r)}} & \\ 2r \sqrt{\frac{n+5r+6}{(n+4r+5)(3+r)}} + (n-4)r \sqrt{\frac{n+5r+7}{(n+4r+6)(3+r)}}. & \end{aligned}$$

Proof. Let $P_n = v_1v_2\dots v_n$ and the r hanging vertices of v_i be $v_i^1, v_i^2, \dots, v_i^r$ ($1 \leq i \leq n$). Let v be a vertex in F_n beside P_n , and the r hanging vertices of v be v^1, v^2, \dots, v^r . By the definition of fourth atom-bond connectivity index, we have

$$\begin{aligned} ABC_4(I_r(F_n)) &= \sum_{i=1}^r \sqrt{\frac{S(v) + S(v^i) - 2}{S(v)S(v^i)}} + \\ \sum_{i=1}^n \sqrt{\frac{S(v) + S(v_i) - 2}{S(v)S(v_i)}} &+ \sum_{i=1}^{n-1} \sqrt{\frac{S(v_i) + S(v_{i+1}) - 2}{S(v_i)S(v_{i+1})}} \\ + \sum_{i=1}^n \sum_{j=1}^r \sqrt{\frac{S(v_i) + S(v_j) - 2}{S(v_i)S(v_j)}} &= \end{aligned}$$

Manuscript received February 09, 2015.

Yun Gao Department of Editorial, Yunnan Normal University, Kunming 650092, China.

Li Liang and Wei Gao School of Information Science and Technology, Yunnan Normal University, Kunming 650500, China

$$\begin{aligned}
 & r\sqrt{\frac{(4n-4)+r(n+2)}{(n+r)((3n-2)+r(n+1))}} \\
 & (2\sqrt{\frac{(4n-1)+r(n+4)}{((3n-2)+r(n+1))(n+3r+3)}} \\
 & +2\sqrt{\frac{(4n+1)+r(n+5)}{((3n-2)+r(n+1))(n+4r+5)}} \\
 & +(n-4)\sqrt{\frac{(4n+2)+r(n+5)}{((3n-2)+r(n+1))(n+4r+6)}} \\
 & (2\sqrt{\frac{2n+8r+9}{(n+4r+5)(n+4r+6)}}+(n-3)\sqrt{\frac{2n+8r+10}{(n+4r+6)(n+4r+6)}}) \\
 & + (2r\sqrt{\frac{n+4r+3}{(n+3r+3)(2+r)}} \\
 & 2r\sqrt{\frac{n+5r+6}{(n+4r+5)(3+r)}}+(n-4)r\sqrt{\frac{n+5r+7}{(n+4r+6)(3+r)}}) \\
 . \quad \square
 \end{aligned}$$

Corollary1.

$$\begin{aligned}
 ABC_4(F_n) &= 2\sqrt{\frac{4n-1}{(3n-2)(n+3)}} \\
 &+ 2\sqrt{\frac{4n+1}{(3n-2)(n+5)}}+(n-4)\sqrt{\frac{4n+2}{(3n-2)(n+6)}} \\
 & 2\sqrt{\frac{2n+9}{(n+5)(n+6)}}+\frac{n-3}{n+6}\sqrt{2n+10}.
 \end{aligned}$$

Theorem 2.

$$\begin{aligned}
 ABC_4(I_r(W_n)) &= r\sqrt{\frac{r(n+2)+4n-2}{(r(n+1)+3n)(n+r)}} \\
 & n\sqrt{\frac{r(n+5)+4n+4}{(r(n+1)+3n)(n+4r+6)}} \\
 & \frac{n}{n+4r+6}\sqrt{2n+8r+10}+nr\sqrt{\frac{n+5r+7}{(n+4r+6)(r+3)}}.
 \end{aligned}$$

Proof. Let $C_n=v_1v_2\dots v_n$ and $v_i^1, v_i^2, \dots, v_i^r$ be the r hanging vertices of v_i ($1 \leq i \leq n$). Let v be a vertex in W_n beside C_n , and v^1, v^2, \dots, v^r be the r hanging vertices of v . According to the definition of fourth atom-bond connectivity index, we get

$$\begin{aligned}
 ABC_4(I_r(W_n)) &= \sum_{i=1}^r \sqrt{\frac{S(v)+S(v^i)-2}{S(v)S(v^i)}} \\
 & \sum_{i=1}^n \sqrt{\frac{S(v)+S(v_i)-2}{S(v)S(v_i)}} + \sum_{i=1}^n \sqrt{\frac{S(v_i)+S(v_{i+1})-2}{S(v_i)S(v_{i+1})}} \\
 & + \sum_{i=1}^n \sum_{j=1}^r \sqrt{\frac{S(v_i)+S(v_i^j)-2}{S(v_i)S(v_i^j)}} \\
 & = r\sqrt{\frac{r(n+2)+4n-2}{(r(n+1)+3n)(n+r)}}
 \end{aligned}$$

$$\begin{aligned}
 & + n\sqrt{\frac{r(n+5)+4n+4}{(r(n+1)+3n)(n+4r+6)}} \\
 & n\sqrt{\frac{2n+8r+10}{(n+4r+6)(n+4r+6)}}+nr\sqrt{\frac{n+5r+7}{(n+4r+6)(r+3)}}.
 \end{aligned}$$

\square

Corollary2.

$$ABC_4(W_n)=n\sqrt{\frac{4n+4}{3n(n+6)}}+\frac{n}{n+6}\sqrt{2n+10}.$$

Theorem3.

$$\begin{aligned}
 ABC_4(I_r(\tilde{F}_n)) &= \\
 & r\sqrt{\frac{(4n-4)+r(n+2)}{(n+r)((3n-2)+r(n+1))}} \\
 & 2\sqrt{\frac{(4n-2)+r(n+4)}{((3n-2)+r(n+1))(n+3r+2)}} \\
 & +(n-2)\sqrt{\frac{4n+r(n+5)}{((3n-2)+r(n+1))(n+4r+4)}} \\
 & 2r\sqrt{\frac{n+4r+2}{(n+3r+2)(2+r)}}+(n-2)r\sqrt{\frac{n+5r+5}{(n+4r+4)(r+3)}} \\
 & 2\sqrt{\frac{n+6r+5}{(n+3r+2)(3r+5)}} \\
 & +2(n-3)\sqrt{\frac{n+7r+8}{(n+4r+4)(3r+6)}}+2\sqrt{\frac{n+7r+7}{(n+4r+4)(3r+5)}} \\
 & +2r\sqrt{\frac{4r+5}{(3r+5)(r+2)}}+(n-2)r\sqrt{\frac{4r+6}{(3r+6)(r+2)}}.
 \end{aligned}$$

Proof. Let $P_n=v_1v_2\dots v_n$ and $v_{i,i+1}$ be the adding vertex between v_i and v_{i+1} . Let $v_i^1, v_i^2, \dots, v_i^r$ be the r hanging vertices of v_i ($1 \leq i \leq n$). Let $v_{i,i+1}^1, v_{i,i+1}^2, \dots, v_{i,i+1}^r$ be the r hanging vertices of $v_{i,i+1}$ ($1 \leq i \leq n-1$). Let v be a vertex in F_n beside P_n , and the r hanging vertices of v be v^1, v^2, \dots, v^r . By virtue of the definition of fourth atom-bond connectivity index, we obtain

$$\begin{aligned}
 ABC_4(I_r(\tilde{F}_n)) &= \sum_{i=1}^r \sqrt{\frac{S(v)+S(v^i)-2}{S(v)S(v^i)}} \\
 & \sum_{i=1}^n \sqrt{\frac{S(v)+S(v_i)-2}{S(v)S(v_i)}} + \sum_{i=1}^n \sum_{j=1}^r \sqrt{\frac{S(v_i)+S(v_i^j)-2}{S(v_i)S(v_i^j)}} \\
 & + \sum_{i=1}^{n-1} \sqrt{\frac{S(v_i)+S(v_{i,i+1})-2}{S(v_i)S(v_{i,i+1})}} + \sum_{i=1}^{n-1} \sqrt{\frac{S(v_{i,i+1})+S(v_{i+1})-2}{S(v_{i,i+1})S(v_{i+1})}} \\
 & + \sum_{i=1}^{n-1} \sum_{j=1}^r \sqrt{\frac{S(v_{i,i+1})+S(v_{i,i+1}^j)-2}{S(v_{i,i+1})S(v_{i,i+1}^j)}}
 \end{aligned}$$

$$\begin{aligned}
 &= r \sqrt{\frac{(4n-4)+r(n+2)}{(n+r)((3n-2)+r(n+1))}} + \\
 &\quad (2\sqrt{\frac{(4n-2)+r(n+4)}{((3n-2)+r(n+1))(n+3r+2)}} \\
 &\quad +(n-2)\sqrt{\frac{4n+r(n+5)}{((3n-2)+r(n+1))(n+4r+4)}}) \\
 &\quad +(2r\sqrt{\frac{n+4r+2}{(n+3r+2)(2+r)}} + (n-2)r\sqrt{\frac{n+5r+5}{(n+4r+4)(r+3)}}) + \\
 &\quad (\sqrt{\frac{n+6r+5}{(n+3r+2)(3r+5)}} \\
 &\quad +(n-3)\sqrt{\frac{n+7r+8}{(n+4r+4)(3r+6)}} + \sqrt{\frac{n+7r+7}{(n+4r+4)(3r+5)}}) \\
 &\quad + (\sqrt{\frac{n+6r+5}{(n+3r+2)(3r+5)}} \\
 &\quad +(n-3)\sqrt{\frac{n+7r+8}{(n+4r+4)(3r+6)}} + \sqrt{\frac{n+7r+7}{(n+4r+4)(3r+5)}}) + \\
 &\quad (2r\sqrt{\frac{4r+5}{(3r+5)(r+2)}} + (n-2)r\sqrt{\frac{4r+6}{(3r+6)(r+2)}}). \square
 \end{aligned}$$

Corollary3.

$$\begin{aligned}
 ABC_4(\tilde{F}_n) &= 2\sqrt{\frac{4n-2}{(3n-2)(n+2)}} \\
 &+(n-2)\sqrt{\frac{4n}{(3n-2)(n+4)}} + 2\sqrt{\frac{n+5}{5(n+2)}} \\
 &+ 2(n-3)\sqrt{\frac{n+8}{6(n+4)}} + 2\sqrt{\frac{n+7}{5(n+4)}}.
 \end{aligned}$$

Theorem4.

$$\begin{aligned}
 ABC_4(I_r(\tilde{W}_n)) &= r\sqrt{\frac{r(n+2)+4n-2}{(r(n+1)+3n)(n+r)}} \\
 &+ n\sqrt{\frac{r(n+5)+4n+4}{(r(n+1)+3n)(n+4r+6)}} \\
 &+ nr\sqrt{\frac{n+5r+7}{(n+4r+6)(r+3)}} + 2n\sqrt{\frac{n+7r+10}{(n+4r+6)(3r+6)}} + \\
 &+ nr\sqrt{\frac{4r+6}{(3r+6)(r+2)}}.
 \end{aligned}$$

Proof. Let $C_n=v_1v_2\dots v_n$ and v be a vertex in W_n beside C_n , $v_{i,i+1}$ be the adding vertex between v_i and v_{i+1} . Let v^1, v^2, \dots, v^r be the r hanging vertices of v and $v_i^1, v_i^2, \dots, v_i^r$ be the r hanging vertices of v_i ($1 \leq i \leq n$). Let $v_{n,n+1}=v_{1,n}$ and $v_{i,i+1}^1, v_{i,i+1}^2, \dots, v_{i,i+1}^r$ be the r hanging vertices of $v_{i,i+1}$ ($1 \leq i \leq n$). In view of the definition of fourth atom-bond connectivity index, we deduce

$$\begin{aligned}
 ABC_4(I_r(\tilde{W}_n)) &= \sum_{i=1}^r \sqrt{\frac{S(v)+S(v^i)-2}{S(v)S(v^i)}} + \\
 &\quad \sum_{i=1}^n \sqrt{\frac{S(v)+S(v_i)-2}{S(v)S(v_i)}} + \sum_{i=1}^n \sum_{j=1}^r \sqrt{\frac{S(v_i)+S(v_j)-2}{S(v_i)S(v_j)}} \\
 &\quad + \sum_{i=1}^n \sqrt{\frac{S(v_i)+S(v_{i,i+1})-2}{S(v_i)S(v_{i,i+1})}} + \sum_{i=1}^n \sqrt{\frac{S(v_{i,i+1})+S(v_{i+1})-2}{S(v_{i,i+1})S(v_{i+1})}} \\
 &\quad + \sum_{i=1}^n \sum_{j=1}^r \sqrt{\frac{S(v_{i,i+1})+S(v_{i,i+1}^j)-2}{S(v_{i,i+1})S(v_{i,i+1}^j)}} \\
 &= r\sqrt{\frac{r(n+2)+4n-2}{(r(n+1)+3n)(n+r)}} + \\
 &\quad n\sqrt{\frac{r(n+5)+4n+4}{(r(n+1)+3n)(n+4r+6)}} + \\
 &\quad nr\sqrt{\frac{n+5r+7}{(n+4r+6)(r+3)}} + n\sqrt{\frac{n+7r+10}{(n+4r+6)(3r+6)}} \\
 &\quad + n\sqrt{\frac{n+7r+10}{(n+4r+6)(3r+6)}} + nr\sqrt{\frac{4r+6}{(3r+6)(r+2)}}. \square
 \end{aligned}$$

Corollary4. $ABC_4(\tilde{W}_n) = n\sqrt{\frac{4n+4}{3n(n+6)}}$

$$+ 2n\sqrt{\frac{n+10}{6(n+6)}}.$$

III GENERAL FIFTH GEOMETRIC-ARITHMETIC INDEX

The terminologies for these special molecular graphs similar as Theorem 1- Theorem 4.

Theorem5.

$$\begin{aligned}
 GA_5^k(I_r(F_n)) &= \\
 &r\left(\frac{2\sqrt{(n+r)((3n-2)+r(n+1))}}{(4n-2)+r(n+2)}\right)^k + \\
 &2\left(\frac{2\sqrt{((3n-2)+r(n+1))(n+3r+3)}}{(4n+1)+r(n+4)}\right)^k + \\
 &+ 2\left(\frac{2\sqrt{((3n-2)+r(n+1))(n+3r+3)}}{(4n+3)+r(n+5)}\right)^k + (n-4)\left(\frac{2\sqrt{((3n-2)+r(n+1))(n+4r+6)}}{(4n+4)+r(n+5)}\right)^k + \\
 &+ 2\left(\frac{2\sqrt{(n+4r+5)(n+4r+6)}}{2n+8r+11}\right)^k + (n-3)\left(\frac{2\sqrt{(n+4r+6)(n+4r+6)}}{2n+8r+12}\right)^k + \\
 &+ 2r\left(\frac{2\sqrt{(n+3r+3)(2+r)}}{n+4r+5}\right)^k + \\
 &+ 2r\left(\frac{2\sqrt{(n+4r+5)(3+r)}}{n+5r+8}\right)^k + (n-4)r\left(\frac{2\sqrt{(n+4r+6)(3+r)}}{n+5r+9}\right)^k.
 \end{aligned}$$

Proof. By the definition of general fifth geometric-arithmetic index, we have

$$\begin{aligned}
 GA_5^k(I_r(F_n)) &= \sum_{i=1}^r \left(\frac{2\sqrt{S(v)S(v^i)}}{S(v)+S(v^i)} \right)^k + r \left(\frac{2\sqrt{(r(n+1)+3n)(n+r)}}{r(n+2)+4n} \right)^k + \\
 &\quad \sum_{i=1}^n \left(\frac{2\sqrt{S(v)S(v_i)}}{S(v)+S(v_i)} \right)^k + \sum_{i=1}^{n-1} \left(\frac{2\sqrt{S(v_i)S(v_{i+1})}}{S(v_i)+S(v_{i+1})} \right)^k + n \left(\frac{2\sqrt{(r(n+1)+3n)(n+4r+6)}}{r(n+5)+4n+6} \right)^k + \\
 &\quad \sum_{i=1}^n \sum_{j=1}^r \left(\frac{2\sqrt{S(v_i)S(v_i^j)}}{S(v_i)+S(v_i^j)} \right)^k + n \left(\frac{2\sqrt{(n+4r+6)(n+4r+6)}}{2n+8r+12} \right)^k + \\
 &= r \left(\frac{2\sqrt{(n+r)((3n-2)+r(n+1))}}{(4n-2)+r(n+2)} \right)^k + nr \left(\frac{2\sqrt{(n+4r+6)(r+3)}}{n+5r+9} \right)^k. \\
 &\quad \square
 \end{aligned}$$

$$\begin{aligned}
 \text{Corollary 5. } GA_5^k(F_n) &= \\
 &2 \left(\frac{2\sqrt{(3n-2)(n+3)}}{4n+1} \right)^k + 2r \left(\frac{2\sqrt{(n+4r+5)(n+4r+6)}}{n+5r+8} \right)^k + (n-4)r \left(\frac{2\sqrt{(n+4r+6)(3+r)}}{n+5r+9} \right)^k. \\
 &\quad \square
 \end{aligned}$$

$$\begin{aligned}
 \text{Theorem 6. } GA_5^k(I_r(W_n)) &= \\
 &r \left(\frac{2\sqrt{(r(n+1)+3n)(n+r)}}{r(n+2)+4n} \right)^k + n \left(\frac{2\sqrt{(r(n+1)+3n)(n+4r+6)}}{r(n+5)+4n+6} \right)^k + nr \left(\frac{2\sqrt{(n+4r+6)(r+3)}}{n+5r+9} \right)^k.
 \end{aligned}$$

Proof. By the definition of general fifth geometric-arithmetic index, we have

$$\begin{aligned}
 GA_5^k(I_r(W_n)) &= \sum_{i=1}^r \left(\frac{2\sqrt{S(v)S(v^i)}}{S(v)+S(v^i)} \right)^k + \\
 &\quad \sum_{i=1}^n \left(\frac{2\sqrt{S(v)S(v_i)}}{S(v)+S(v_i)} \right)^k + \sum_{i=1}^n \left(\frac{2\sqrt{S(v_i)S(v_{i+1})}}{S(v_i)+S(v_{i+1})} \right)^k + \\
 &\quad \sum_{i=1}^n \sum_{j=1}^r \left(\frac{2\sqrt{S(v_i)S(v_i^j)}}{S(v_i)+S(v_i^j)} \right)^k
 \end{aligned}$$

$$\begin{aligned}
 &= r \left(\frac{2\sqrt{(r(n+1)+3n)(n+r)}}{r(n+2)+4n} \right)^k + \\
 &\quad n \left(\frac{2\sqrt{(r(n+1)+3n)(n+4r+6)}}{r(n+5)+4n+6} \right)^k + \\
 &\quad n \left(\frac{2\sqrt{(n+4r+6)(n+4r+6)}}{2n+8r+12} \right)^k + \\
 &+ nr \left(\frac{2\sqrt{(n+4r+6)(r+3)}}{n+5r+9} \right)^k. \\
 &\quad \square
 \end{aligned}$$

$$\text{Corollary 6. } GA_5^k(W_n) = n \left(\frac{2\sqrt{3n(n+6)}}{4n+6} \right)^k + n.$$

$$\begin{aligned}
 \text{Theorem 7. } GA_5^k(I_r(\tilde{F}_n)) &= \\
 &r \left(\frac{2\sqrt{(n+r)((3n-2)+r(n+1))}}{(4n-2)+r(n+2)} \right)^k + \\
 &2 \left(\frac{2\sqrt{((3n-2)+r(n+1))(n+3r+2)}}{4n+r(n+4)} \right)^k + \\
 &+(n-2) \left(\frac{2\sqrt{((3n-2)+r(n+1))(n+4r+4)}}{(4n+2)+r(n+5)} \right)^k + \\
 &2r \left(\frac{2\sqrt{(n+3r+2)(2+r)}}{n+4r+4} \right)^k + \\
 &+(n-2)r \left(\frac{2\sqrt{(n+4r+4)(r+3)}}{n+5r+7} \right)^k + \\
 & \left(\frac{2\sqrt{(n+3r+2)(3r+5)}}{n+6r+7} \right)^k + \\
 &+(n-3) \left(\frac{2\sqrt{(n+4r+4)(3r+6)}}{n+7r+10} \right)^k + \\
 &+\left(\frac{2\sqrt{(n+4r+4)(3r+5)}}{n+7r+9} \right)^k + \\
 & \left(\frac{2\sqrt{(n+3r+2)(3r+5)}}{n+6r+7} \right)^k + \\
 &+(n-3) \left(\frac{2\sqrt{(n+4r+4)(3r+6)}}{n+7r+10} \right)^k + \\
 &+\left(\frac{2\sqrt{(n+4r+4)(3r+5)}}{n+7r+9} \right)^k + \\
 &2r \left(\frac{2\sqrt{(3r+5)(r+2)}}{4r+7} \right)^k + (n-2)r \left(\frac{2\sqrt{(3r+6)(r+2)}}{4r+8} \right)^k.
 \end{aligned}$$

Proof. By virtue of the definition of general fifth geometric-arithmetic index, we get

$$\begin{aligned}
 GA_5^k(I_r(\tilde{F}_n)) &= \sum_{i=1}^r \left(\frac{2\sqrt{S(v)S(v^i)}}{S(v)+S(v^i)} \right)^k + \\
 &\quad \sum_{i=1}^n \left(\frac{2\sqrt{S(v)S(v_i)}}{S(v)+S(v_i)} \right)^k + \sum_{i=1}^n \sum_{j=1}^r \left(\frac{2\sqrt{S(v_i)S(v_i^j)}}{S(v_i)+S(v_i^j)} \right)^k +
 \end{aligned}$$

$$\begin{aligned}
 & \sum_{i=1}^{n-1} \left(\frac{2\sqrt{S(v_i)S(v_{i,i+1})}}{S(v_i)+S(v_{i,i+1})} \right)^k + \sum_{i=1}^{n-1} \left(\frac{2\sqrt{S(v_{i,i+1})S(v_{i+1})}}{S(v_{i,i+1})+S(v_{i+1})} \right)^k + \\
 & \sum_{i=1}^{n-1} \sum_{j=1}^r \left(\frac{2\sqrt{S(v_{i,i+1})S(v_{i,i+1}^j)}}{S(v_{i,i+1})+S(v_{i,i+1}^j)} \right)^k \\
 = & r \left(\frac{2\sqrt{(n+r)((3n-2)+r(n+1))}}{(4n-2)+r(n+2)} \right)^k \\
 & (2 \left(\frac{2\sqrt{((3n-2)+r(n+1))(n+3r+2)}}{4n+r(n+4)} \right)^k \\
 & +(n-2) \left(\frac{2\sqrt{((3n-2)+r(n+1))(n+4r+4)}}{(4n+2)+r(n+5)} \right)^k) \\
 & (2r \left(\frac{2\sqrt{(n+3r+2)(2+r)}}{n+4r+4} \right)^k \\
 & +(n-2)r \left(\frac{2\sqrt{(n+4r+4)(r+3)}}{n+5r+7} \right)^k) \\
 & ((\left(\frac{2\sqrt{(n+3r+2)(3r+5)}}{n+6r+7} \right)^k \\
 & +(n-3) \left(\frac{2\sqrt{(n+4r+4)(3r+6)}}{n+7r+10} \right)^k + \left(\frac{2\sqrt{(n+4r+4)(3r+5)}}{n+7r+9} \right)^k) \\
 & + ((\left(\frac{2\sqrt{(n+3r+2)(3r+5)}}{n+6r+7} \right)^k \\
 & +(n-3) \left(\frac{2\sqrt{(n+4r+4)(3r+6)}}{n+7r+10} \right)^k + \left(\frac{2\sqrt{(n+4r+4)(3r+5)}}{n+7r+9} \right)^k) \\
 & + (2r \left(\frac{2\sqrt{(3r+5)(r+2)}}{4r+7} \right)^k \\
 & +(n-2)r \left(\frac{2\sqrt{(3r+6)(r+2)}}{4r+8} \right)^k) \\
 \end{aligned}$$

□

Corollary7.

$$\begin{aligned}
 & GA_5^k(\tilde{F}_n) \\
 & 2 \left(\frac{2\sqrt{(3n-2)(n+2)}}{4n} \right)^k + (n-2) \left(\frac{2\sqrt{(3n-2)(n+4)}}{4n+2} \right)^k + \\
 & \left(\frac{2\sqrt{5(n+2)}}{n+7} \right)^k \\
 & + 2(n-3) \left(\frac{2\sqrt{6(n+4)}}{n+10} \right)^k + 2 \left(\frac{2\sqrt{5(n+4)}}{n+9} \right)^k \\
 & \left(\frac{2\sqrt{5(n+2)}}{n+7} \right)^k.
 \end{aligned}$$

Theorem8.

$$\begin{aligned}
 & GA_5^k(I_r(\tilde{W}_n)) \\
 & r \left(\frac{2\sqrt{(r(n+1)+3n)(n+r)}}{r(n+2)+4n} \right)^k \\
 & n \left(\frac{2\sqrt{(r(n+1)+3n)(n+4r+6)}}{r(n+5)+4n+6} \right)^k \\
 & nr \left(\frac{2\sqrt{(n+4r+6)(r+3)}}{n+5r+9} \right)^k
 \end{aligned}$$

$$\begin{aligned}
 & + 2n \left(\frac{2\sqrt{(n+4r+6)(3r+6)}}{n+7r+12} \right)^k + \\
 & nr \left(\frac{2\sqrt{(3r+6)(r+2)}}{4r+8} \right)^k. \\
 \text{Proof.} \quad & \text{In view of the definition of general fifth geometric-arithmetic index, we deduce} \\
 GA_5^k(I_r(\tilde{W}_n)) & = \sum_{i=1}^r \left(\frac{2\sqrt{S(v)S(v^i)}}{S(v)+S(v^i)} \right)^k + \\
 & \sum_{i=1}^n \left(\frac{2\sqrt{S(v)S(v_i)}}{S(v)+S(v_i)} \right)^k + \sum_{i=1}^n \sum_{j=1}^r \left(\frac{2\sqrt{S(v_i)S(v_i^j)}}{S(v_i)+S(v_i^j)} \right)^k + \\
 & \sum_{i=1}^n \left(\frac{2\sqrt{S(v_i)S(v_{i,i+1})}}{S(v_i)+S(v_{i,i+1})} \right)^k + \sum_{i=1}^n \left(\frac{2\sqrt{S(v_{i,i+1})S(v_{i+1})}}{S(v_{i,i+1})+S(v_{i+1})} \right)^k + \\
 & \sum_{i=1}^n \sum_{j=1}^r \left(\frac{2\sqrt{S(v_{i,i+1})S(v_{i,i+1}^j)}}{S(v_{i,i+1})+S(v_{i,i+1}^j)} \right)^k \\
 = & r \left(\frac{2\sqrt{(r(n+1)+3n)(n+r)}}{r(n+2)+4n} \right)^k + \\
 & n \left(\frac{2\sqrt{(r(n+1)+3n)(n+4r+6)}}{r(n+5)+4n+6} \right)^k + \\
 & nr \left(\frac{2\sqrt{(n+4r+6)(r+3)}}{n+5r+9} \right)^k \\
 & + n \left(\frac{2\sqrt{(n+4r+6)(3r+6)}}{n+7r+12} \right)^k + \\
 & nr \left(\frac{2\sqrt{(3r+6)(r+2)}}{4r+8} \right)^k. \square
 \end{aligned}$$

$$\begin{aligned}
 & \text{Corollary8.} \quad GA_5^k(\tilde{W}_n) \\
 & n \left(\frac{2\sqrt{3n(n+6)}}{4n+6} \right)^k + 2n \left(\frac{2\sqrt{6(n+6)}}{n+12} \right)^k.
 \end{aligned}$$

IV FIFTH GEOMETRIC-ARITHMETIC INDEX

By taking $k=1$ in Theorem 5-8, we get the following conclusions on fifth geometric-arithmetic index.

Theorem9.

$$\begin{aligned}
 GA_5(I_r(F_n)) & = \frac{2r\sqrt{(n+r)((3n-2)+r(n+1))}}{(4n-2)+r(n+2)} + \\
 & 4\sqrt{((3n-2)+r(n+1))(n+3r+3)} \\
 & \quad (4n+1)+r(n+4) \\
 & + \frac{4\sqrt{((3n-2)+r(n+1))(n+4r+5)}}{(4n+3)+r(n+5)} + \frac{2(n-4)\sqrt{((3n-2)+r(n+1))(n+4r+6)}}{(4n+4)+r(n+5)} \\
 & + \frac{4\sqrt{(n+4r+5)(n+4r+6)}}{2n+8r+11} \\
 & + \frac{2(n-3)\sqrt{(n+4r+6)(n+4r+6)}}{2n+8r+12}
 \end{aligned}$$

$$\frac{4r\sqrt{(n+3r+3)(2+r)}}{n+4r+5} + \frac{4r\sqrt{(n+4r+5)(3+r)}}{n+5r+8} + \frac{2(n-4)r\sqrt{(n+4r+6)(3+r)}}{n+5r+9}.$$

Corollary9.

$$GA_5(F_n) = \frac{4\sqrt{(3n-2)(n+3)}}{4n+1} + \frac{4\sqrt{(3n-2)(n+5)}}{4n+3} + \frac{2(n-4)\sqrt{(3n-2)(n+6)}}{4n+4} + \frac{4\sqrt{(n+5)(n+6)}}{2n+11} + \frac{2(n-3)(n+6)}{2n+12}.$$

Theorem

$$GA_5(I_r(W_n)) = \frac{2r\sqrt{(r(n+1)+3n)(n+r)}}{r(n+2)+4n} + \frac{2n\sqrt{(r(n+1)+3n)(n+4r+6)}}{r(n+5)+4n+6} + \frac{2nr\sqrt{(n+4r+6)(r+3)}}{n+5r+9}.$$

$$\text{Corollary 10. } GA_5(W_n) = \frac{2n\sqrt{3n(n+6)}}{4n+6} + n.$$

Theorem

$$GA_5(I_r(\tilde{F}_n)) = \frac{2r\sqrt{(n+r)((3n-2)+r(n+1))}}{(4n-2)+r(n+2)} + \frac{4\sqrt{((3n-2)+r(n+1))(n+3r+2)}}{4n+r(n+4)} + \frac{2(n-2)\sqrt{((3n-2)+r(n+1))(n+4r+4)}}{(4n+2)+r(n+5)} + \frac{4r\sqrt{(n+3r+2)(2+r)}}{n+4r+4} + \frac{2(n-2)r\sqrt{(n+4r+4)(r+3)}}{n+5r+7} + \frac{2\sqrt{(n+3r+2)(3r+5)}}{n+6r+7}$$

$$+ \frac{2(n-3)\sqrt{(n+4r+4)(3r+6)}}{n+7r+10} + \frac{2\sqrt{(n+4r+4)(3r+5)}}{n+7r+9} + \frac{2\sqrt{(n+3r+2)(3r+5)}}{n+6r+7} + \frac{2(n-3)\sqrt{(n+4r+4)(3r+6)}}{n+7r+10} + \frac{2\sqrt{(n+4r+4)(3r+5)}}{n+7r+9} + \frac{4r\sqrt{(3r+5)(r+2)}}{4r+7} + \frac{2(n-2)r\sqrt{(3r+6)(r+2)}}{4r+8}.$$

Corollary11.

$$GA_5(\tilde{F}_n) = \frac{\sqrt{(3n-2)(n+2)}}{n} + \frac{(n-2)\sqrt{(3n-2)(n+4)}}{2n+1}$$

$$\frac{2\sqrt{5(n+2)}}{n+7} + \frac{2(n-3)\sqrt{6(n+4)}}{n+10} + \frac{2\sqrt{5(n+4)}}{n+9} + \frac{2\sqrt{5(n+2)}}{n+7} + \frac{2(n-3)\sqrt{6(n+4)}}{n+10} + \frac{2\sqrt{5(n+4)}}{n+9}.$$

Theorem12.

$$GA_5(I_r(\tilde{W}_n)) = \frac{2r\sqrt{(r(n+1)+3n)(n+r)}}{r(n+2)+4n} + \frac{2n\sqrt{(r(n+1)+3n)(n+4r+6)}}{r(n+5)+4n+6} + \frac{2nr\sqrt{(n+4r+6)(r+3)}}{n+5r+9} + \frac{4n\sqrt{(n+4r+6)(3r+6)}}{n+7r+12} + \frac{2nr\sqrt{(3r+6)(r+2)}}{4r+8}.$$

$$\text{Corollary 12. } GA_5(\tilde{W}_n) = \frac{2n\sqrt{3n(n+6)}}{4n+6} + \frac{2n\sqrt{6(n+6)}}{n+12} + \frac{2n\sqrt{6(n+6)}}{n+12}.$$

10.

11.

V. CONCLUSION

In this paper, we determine the fourth atom bond connectivity index and fifth geometric-arithmetic index of fan molecular graph, wheel molecular graph, gear fan molecular graph, gear wheel molecular graph, and their r -corona molecular graphs. Furthermore, the general version of fifth geometric-arithmetic index is discussed.

ACKNOWLEDGEMENTS

We thank the reviewers for their constructive comments in improving the quality of this paper. This work was supported in part by NSFC (no. 11401519). We also would like to thank the anonymous referees for providing us with constructive comments and suggestions.

REFERENCES

- [1] L. Yan, Y. Li, W. Gao, J. S. Li, On the extremal hyper-wiener index of graphs, Journal of Chemical and Pharmaceutical Research, 2014, 6(3): 477-481.
- [2] L. Yan, W. Gao, J. S. Li, General harmonic index and general sum connectivity index of polyomino chains and nanotubes, Journal of Computational and Theoretical Nanoscience, In press.
- [3] W. Gao, L. Liang, Y. Gao, Some results on wiener related index and shultz index of molecular graphs, Energy Education Science and Technology: Part A, 2014, 32(6): 8961-8970.
- [4] W. Gao, L. Liang, Y. Gao, Total eccentricity, adjacent eccentric distance sum and Gutman index of certain special molecular graphs, The BioTechnology: An Indian Journal, 2014, 10(9): 3837-3845.
- [5] W. Gao, L. Shi, Wiener index of gear fan graph and gear wheel graph, Asian Journal of Chemistry, 2014, 26(11): 3397-3400.
- [6] W. Gao, W. F. Wang, Second atom-bond connectivity index of special chemical molecular structures, Journal of Chemistry, Volume 2014, Article ID 906254, 8 pages, <http://dx.doi.org/10.1155/2014/906254>.
- [7] W. F. Xi, W. Gao, Geometric-arithmetic index and Zagreb indices of certain special molecular graphs, Journal of Advances in Chemistry, 2014, 10(2): 2254-2261.
- [8] W. F. Xi, W. Gao, λ -Modified extremal hyper-Wiener index of

- molecular graphs, *Journal of Applied Computer Science & Mathematics*, 2014, 18 (8): 43-46.
- [9] W. F. Xi, W. Gao, Y. Li, Three indices calculation of certain crown molecular graphs, *Journal of Advances in Mathematics*, 2014, 9(6): 2696-2304.
- [10] Y. Gao, W. Gao, L. Liang, Revised Szeged index and revised edge Szeged index of certain special molecular graphs, *International Journal of Applied Physics and Mathematics*, 2014, 4(6): 417-425.
- [11] J. A. Bondy, U. S. R. Murty, *Graph Theory*, Springer, Berlin, 2008.
- [12] M. Ghorbani, M. A. Hosseini, Computing ABC₄ index of nanostar dendrimers, *Optoelectron. Adv. Mater.-Rapid Commun.* 2010, 4(9):1419-1422.
- [13] M. Ghorbani, M. Jalili, Computing a New Topological Index of Nano Structures, *Digest. J. Nanomater. Bios.* 2009, 4(4): 681 - 685.
- [14] M. Ghorbani, M. Ghazi, Computing some topological indices of Triangular Benzenoid, *Digest. J. Nanomater. Bios.* 2010, 5(4): 1107-1111.
- [15] M. R. Farahani, Fourth Atom-Bond Connectivity (ABC4) Index of Nanostructures, *Sci-Afric Journal of Scientific Issues, Research and Essays*, 2014, 2 (12): 567-570.
- [16] A. Graovac, M. Ghorbani, M. A. Hosseini, Computing fifth geometric-arithmetic index for Nanostar Dendrimers, *Journal of Mathematical Nano Science*, 2011, 1 (1): 32-42.
- [17] M. R. Farahani, Fifth geometric-arithmetic index of Polyhex Zigzag TUZC₆[m;n] nanotube and nanotori, *Journal of Advances in Physics*, 2013, 3(1): 191-196.
- [18] M. R. Farahani, Computing GA₅ index of armchair polyhex nanotube, *LE MATEMATICHE*, 2014, Vol. LXIX – Fasc. II: 69–76.

Yun Gao was born in Yunnan Province, China, on December 10, 1964. He studied in the Department of Physics of Yunnan Normal University from 1980 to 1984 and got his bachelor degree on physics. And then he worked as an editor and a physics lecturer in the Journal of Yunnan Normal University till now. During his years, he was always engaged in studying the computational physics and other related areas, such as condensed matter physics and computer science. He had published more than 30 scientific papers in home and abroad. Now, as the vice editor in chief of Yunnan Normal University and a researcher, his interests are mainly in computational physics and computing method.

Li Liang was born in 1965. As a professor of Yunnan Normal University, his interests are in graph theory, statistical learning theory, information retrieval, and artificial intelligence

Wei Gao was born in the City of Shaoxing, Zhejiang Province, China on February 13, 1981. He got two bachelor degrees on computer science from Zhejiang Industrial University in 2004 and mathematics education form College of Zhejiang Education in 2006. Then, he enrolled in the Department of Computer Science and Information Technology, Yunnan Normal University, and got his master degree there in 2009. In 2012, he got PhD degree in the Department of Mathematics, Soochow University, China. Now, he acts as a lecturer in the Department of Information, Yunnan Normal University. As a researcher in computer science and mathematics, his interests are covered two disciplines: graph theory, statistical learning theory, information retrieval, and artificial intelligence