

Some Properties of Semi-continuous, Pre-continuous and α -continuous Mappings

Yasser A. Abou-Elwan

Abstract— This paper investigates some new characteristics of semi-continuous, pre-continuous and α -continuous mappings. We provides two theorems that are equivalent to the definitions of pre-continuous and M-semi-continuous mappings. A condition has been proposed, which makes the injective mapping pre-open. We have proved that the domain of the injective α -continuous mapping with closed graph is Housdorff space. In addition, more other conditions put on the α -continuous mapping, which make its graph closed.

Index Terms—closed graph, semi-continuous mappings, pre-continuous mapping, α -continuous mapping, α -open mapping, M-semi-continuous mapping

I. INTRODUCTION

A subset A of the space X is called a semi-open [5] (resp. α -set [7], pre-open [1], β -open [2], regular open[8]) $A \subset A^{s-}$ (resp. $A \subset A^{s+}$, $A \subset A^{-s}$, $A \subset A^{-s+}$, $A = A^{-s}$). The complement of a semi-open (resp. α -set, pre-open, β -set, regular open) set, is called a semi-closed[5] (resp. α -closed [6], pre-closed[1], β -closed[2], regular closed[7]). The family of all semi-open (resp. α -set, pre-open, β -open, regular open) sets of a space X will be denoted by $SO(X)$ (resp. $\alpha(X)$, $PO(X)$, $\beta O(X)$, $RO(X)$).

A mapping $f : X \rightarrow Y$ is called semi-continuous [5] (resp. α -continuous [6], pre-continuous [1], and β -continuous [4]) if the inverse image of every open set in Y is semi-open (resp. α -set, pre-open, β -open) in X .

Theorem 1.1. [5]. Let $f : X \rightarrow Y$ be a mapping, then the following statements are equivalent:

- i) f is β -continuous.
- ii) For every $x \in X$ and every open set $V \subset Y$ containing $f(x)$, there exists a β -open set $W \subset X$ containing x such that $f(W) \subset V$.
- iii) The inverse image of each closed set in Y is β -closed in X .
- iv) $(f^{-1}(B))^{o-o} \subset f^{-1}(\overline{B})$, for every $B \subset Y$.
- v) $f(A^{o-o}) \subset \overline{f(A)}$, for every $A \subset X$.

Theorem 1.2. [1]. Let $f : X \rightarrow Y$ be a mapping, then the following statements are equivalent:

- i) f is β -open.
- ii) For every $x \in X$ and every neighborhood U of x , there exists a β -open set $W \subset Y$ containing $f(x)$ such that $W \subset f(U)$.
- iii) $f^{-1}(B^{o-o}) \subset f^{-1}(\overline{B})$, for every $B \subset Y$.
- iv) If f is bijective, then $(f(A))^{o-o} \subset \overline{f(A)}$, for every $A \subset X$.

Definition 1.1. [3],[5] A mapping $f : X \rightarrow Y$ is said to have a closed graph , if its graph $G(f) = \{(x, y) : y = f(x), x \in X\}$ in the product space $X \times Y$ is a closed set . Equivalently ,

$G(f)$ is a closed subset of $X \times Y$, if and only of for each $x \in X$, and $y \neq f(x)$, there exist open sets U and V containing x and y respectively such that $f(U) \cap V = \emptyset$.

II. SEMI-CONTINUOUS, PRE-CONTINUOUS AND α -CONTINUOUS MAPPINGS.

Theorem 2.1. A mapping $f : X \rightarrow Y$ is pre-continuous iff $f(\overline{U}) \subset (f(U))^{-}$, for every open set $U \subset X$.

Proof. Let f be a pre-continuous, then by theorem 1.1., $f(U^{s-}) \subset (f(U))^{-}$, for every U , since $U \subset X$, since $U \subset X$ is open, then $f(\overline{U}) \subset (f(U))^{-}$.

Conversely, let $V \subset Y$ be open, $W = Y - V$, and let $U = (f^{-1}(W))^{s-}$ be an open subset of X , then

$$f(f^{-1}(W))^{s-} \subset (f((f^{-1}(W))^{s-}))^{-} \subset (f(f^{-1}(W)))^{-} \subset \overline{W} = W$$

. So, $(f^{-1}(W))^{s-} \subset f^{-1}(W)$ and f is pre-continuous .

Theorem 2.2. An injective mapping $f : X \rightarrow Y$ is pre-open iff $f^{-1}(\overline{B}) \subset (f^{-1}(B))^{-}$, for every open set $B \subset Y$.

Proof. Suppose f is pre-open then by Theorem 1.2., $f^{-1}(B^{s-}) \subset (f^{-1}(B))^{-}$. Since $B \subset Y$ is open,

$$f^{-1}(\overline{B}) \subset (f^{-1}(B))^{-}$$

Conversely, let $V \subset X$ be an open set , $W = X - V$ and let $B = (f(W))^{o}$ be an open subset of Y .

Then

$$f^{-1}((f(W))^{o}) \subset (f^{-1}(W))^{s-} \subset (f^{-1}(f(W)))^{-} = \overline{W} = W.$$

Hence $(f(W))^{s-} \subset f(W)$, and so, f is pre-open.

Theorem 2.3. Let $f : X \rightarrow Y$ be an injective α -continuous mapping with closed graph. Then X is a T_2 -space.

Proof. Let $x_1, x_2 \in X$, $x_1 \neq x_2$, then $f(x_1) \neq f(x_2)$. Since $G(f)$ is closed , then there exist open sets U and V containing x_1 and $f(x_2)$ respectively , such that $f(U) \cap V = \emptyset$. Thus $f^{-1}(V) \subset X - U$, and $(f^{-1}(V))^{s-} \subset (X - U)^{s-} = (X - U^{-s})^{s-}$. Since f is α -continuous, $x_2 \in f^{-1}(V) \subset (X - \overline{U})^{s-}$. Therefore, X is a T_2 -space.

Theorem 2.4. Let $f : X \rightarrow Y$ be α -continuous mapping where Y is locally connected T_2 -space. If f and f^{-1} map connected sets into connected sets, then the graph $G(f)$ of f is closed.

Proof. Let $x \in X$, and $y \in f(x)$, $y \in Y$. Since Y is T_2 -space, there exist two disjoint open sets U and V containing y and $f(x)$, respectively. Since Y is locally connected , there exist an

open connected set W such that $f(x) \in W \subset V$. So $W \cap U = \phi$ and $f^{-1}(W) \cap f^{-1}(U) = \phi$. Since f is α -continuous,

$$x \in f^{-1}(W) \subset (f^{-1}(W))^{s-}$$

Now since any point $p \in f^{-1}(U)$ is a limit point of the connected set $f^{-1}(W)$, then $\{p\} \cup f^{-1}(W)$ is connected. But, $f(\{p\} \cup f^{-1}(W))$ has points in each of the two disjoint open sets U and W and so, $f(\{p\} \cup f^{-1}(W))$ is not connected, which is a contradiction to our assumption that f maps connected sets into connected sets. Hence $(f^{-1}(W))^{s-} \cap f^{-1}(U) = \phi$ and $f(f^{-1}(W))^{s-} \cap U = \phi$, therefore $G(f)$ is closed.

Theorem 2.5. Let f be α -continuous surjection, then Y is connected if X is connected.

Proof. Assume Y is not connected and X is connected, then there are two disjoint open sets $V_i \subset Y$, $i \in \{1,2\}$ such that $U_i V_i = Y$ and $\bigcap_i V_i = \phi$. Since f is α -continuous and since α -continuity implies β -continuity, then $f^{-1}(V_i) \subset (f^{-1}(V_i))^{s-} \subset f^{-1}(\bar{V}_i)$. Since V_i is open and closed for every $i \in \{1,2\}$,

$\bigcap_i f^{-1}(V_i) \subset \bigcap_i (f^{-1}(V_i))^{s-} \subset \bigcap_i f^{-1}(\bar{V}_i) \cap f^{-1}(V_i) = \phi$. Hence X is not connected, and this leads to a contradiction which proves that Y is connected.

Theorem 2.6. For a bijective mapping f , f is α -open iff $(f(U))^{s-} \subset f(\bar{U})$, for every $U \subset X$.

Proof. Suppose f is α -open. Let $U \subset X$, then $f(X - \bar{U}) \subset (f(X - \bar{U}))^{s-} \subset (f(X - U))^{s-}$. Since f is bijective, $f(\bar{U}) \supset (f(U))^{s-}$. Conversely, suppose U is an open set of X . Then $f(X - U) = f(X - U) \supset (f(X - U))^{s-}$. Since f is bijective, $f(U) \subset (f(U))^{s-}$, and so, f is α -open.

Definition 2.1.[1]. A mapping $f : X \rightarrow Y$ is called M-semi-continuous if the inverse image of every semi-open set in Y is semi-open in X .

The following theorem gives a new property of a semi-continuous mappings.

Theorem 2.7. Let $f : X \rightarrow Y$ be semi-continuous and $f^{-1}(\bar{V}) \subset (f^{-1}(V))^-$ for every semi-open set $V \subset Y$, then f is M-semi-continuous.

Proof. Let V be semi-open set in X . Since f is semi-continuous, Then $f^{-1}(V) \subset f^{-1}(V^{s-}) \subset (f^{-1}(V^o))^- \subset (f^{-1}(V^*))^s = f^{-1}(V^*)^{s-} \subset (f^{-1}(V))^{s-}$. Hence f is M-semi-continuous.

REFERENCES

[1] Abd El-Monsef, M.E. , . Studies on some pre –topological concepts, vol.11,*pacific, J. Math.*, 1980, pp.339-347.
 [2] Abd El-Monsef, M.E., El-Deeb, S .N., Mahmoud, F.A., On β - open set,vol. 13, *pacific, J. Math.*, 1981, pp.561-569.
 [3] Husain, J., *Topology and Maps* , *Plenum Press, New York*,1977.
 [4] Levine. N., A de composition of continuity in topological spaces, vol.68, *Amer. Math. Monthly*, 1961, pp.44-46 .
 [5] Levine. N. , Semi-open sets and semi continuity in topological spaces, vol.70, *Amer. Math. Monthly*, 1963, pp.36-41
 [6] Mashhour, A. S.; Hasanian, I. A.;El-Deeb, S. N. , α - continuous and α - open mappings, vol.41, *Acta. Math. Hungar*, 1983, pp.213-218.
 [7] Najstad, O., On some classes of nearly open sets, vol.15, *pacific, J. Math.*, 1965, pp.961-970.
 [8] Singal, M. J., Rani, A., Almost-continuous mappings, vol.16, *Yokohama, Math. J.*, 1968, pp.63 -73.