

Applications of Two Dimensional Fractional Fourier-Mellin Transform to Differential Equations

V.D. Sharma, P.B. Deshmukh

Abstract— Integral transforms play wide and important role in mathematical physics, theoretical physics. Fourier-Mellin transform is used in fields like electronics, agriculture, medical etc.

It has applications as registration of images, watermarks, invariant pattern recognition, preprocessing of images.

In this paper we have obtained the differential operator Λ and Λ^* . Using it we have solved the differential equation of the type . Also an application of two-dimensional fractional Fourier-Mellin transform to differential equation is presented.

Index Terms— Two-Dimensional Fractional Fourier-Mellin Transform, Testing Function Space, Generalized function.

I. INTRODUCTION

Invariant content-based image retrieval using a complete set of Fourier-Mellin descriptors works better for database containing isolated objects on a uniform background, it seems well-suited for professional database such as : medical, biology, telecommunication etc [1].

Hyperspectral imagery is playing important role in many fields such as geology, agriculture, environment, military, atmosphere and so on. Image registration with hyperspectral data based on fourier-mellin transform [2]. A visual odometry method that estimates the location and orientation of a robotic rover platform. Visual Odometry Based on the Fourier-Mellin Transform for a Rover Using a Monocular Ground-Facing Camera [3]. RMI, RTMT are basically used in medical for image registration is based on fourier-mellin transform . This transform gives a fast registration and good registration accuracy and avoid trapping in the local optima [4].

Fourier-Mellin transform provides a global method for registering images in a video sequence from which the rotation and translation of the camera motion can be estimated. It has been ability of the gray level image representation for pattern recognition. FMT is used to identify plant leaves at various life stages based on the leaves shape or contour [9]. Milanese, R.,Cherbuliez, M.,Pun, T. was looking for the invariant content of images in multimedia archives using Fourier-Mellin transformation[10].

Motivated by the above work, in our previous work we have generalized two dimensional fractional Fourier-Mellin transform in the distributional sense. Also proved the analyticity theorem, an inversion theorem, convolution theorem, also calculated adjoint operators. [5,6,7,8]. In the present work we have obtained the differential operator Λ and Λ^* . Using it we have solved the differential equation of the

Dr. V. D. Sharma, Assistant professor in the department of Mathematics, Arts, Commerce and Science College, Kiran Nagar, Amravati-444606 (M.S.) India.

P. B. Deshmukh, Assistant professor in the department of Mathematics, IBSS College of Engineering, Amravati 444602.

type $P(\Lambda^*_{x,y,t,q})v = g$. Also an application of two-dimensional fractional Fourier-Mellin transform to differential equation is calculated.

II. TWO-DIMENSIONAL FRACTIONAL FOURIER-MELLIN TRANSFORM

A. Definition of two-dimensional fractional Fourier-Mellin transform:

The two-dimensional fractional Fourier-Mellin transform with parameters α and θ of $f(x, y, t, q)$ denoted by

2DFRFMT{ $f(x, y, t, q)$ } performs a linear operation, given

by the integral transform. 2DFRFMT

$$\{ f(x, y, t, q) \} = F_{\alpha, \theta}(\xi, \eta, \lambda, \chi)$$

$$\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_0^{\infty} \int_0^{\infty} f(x, y, t, q) K_{\alpha, \theta}(x, y, t, q, \xi, \eta, \lambda, \chi) dx dy dt dq \text{----(1)}$$

Where $K_{\alpha, \theta}(x, y, t, q, \xi, \eta, \lambda, \chi)$

$$= \sqrt{\frac{1 - icot\alpha}{2\pi}} e^{\frac{1}{2i\sin\alpha} i(x^2 + y^2 + \xi^2 + \eta^2) \cos\alpha - 2(x\xi + y\eta)}$$

$$e^{\frac{2\pi i \lambda}{t \sin\theta} - 1} e^{\frac{2\pi i \chi}{q \sin\theta} - 1} e^{\frac{\pi i}{t \tan\theta} (t^2 + \lambda^2 + \log^2 t + \log^2 q)}$$

$$= C_{1\alpha} e^{iC_{2\alpha} i(x^2 + y^2 + \xi^2 + \eta^2) \cos\alpha - 2(x\xi + y\eta)}$$

$$e^{iC_{1\theta} i\lambda - 1} e^{iC_{1\theta} i\chi - 1} e^{iC_{2\theta} i(t^2 + \lambda^2 + \log^2 t + \log^2 q)}$$

$$\text{where } C_{1\alpha} = \sqrt{\frac{1 - icot\alpha}{2\pi}}, C_{2\alpha} = \frac{1}{2i\sin\alpha} C_{1\theta} = \frac{2\pi}{\sin\theta},$$

$$C_{2\theta} = \frac{\pi}{\tan\theta} \quad 0 < \alpha < \frac{\pi}{2}, \quad 0 < \theta < \frac{\pi}{2} \text{----(2)}$$

B. The Test Function

An infinitely differentiable complex valued smooth function $\phi(x, y, t, q)$ on R^n belongs to $E(R^n)$, if for each compact set $I \subset S_{a,b}$, $J \subset S_{c,d}$ where

$$S_{a,b} = \{x, y: x, y \in R^n, |x| \leq a, |y| \leq b, a > 0, b > 0\}$$

$$S_{c,d} = \{t, q: t, q \in R^n, |t| \leq c, |q| \leq d, c > 0, d > 0\}$$

$$Y_{E,m,n,k,l}[\phi(x,y,t,q)] = \sup_{x,y,t,q} |D_{x,y,t,q}^{m,n,k,l} \phi(x,y,t,q)| < \infty \text{---(3)}$$

Thus $E(R^n)$ will denote the space of all

$\phi(x, y, t, q) \in E(R^n)$ with compact support contained in

$$S_{a,b} \cap S_{c,d}.$$

Note that the space E is complete and therefore a Fréchet space. Moreover, we say that $f(x, y, t, q)$ is a fractional Fourier-Mellin transformable if it is a member of E .

III. DISTRIBUTIONAL TWO DIMENSIONAL FRACTIONAL FOURIER-MELLIN TRANSFORM (2DFRFMT)

The two dimensional distributional Fractional Fourier -Mellin transform of $f(x, y, t, q) \in E^*(\mathbb{R}^n)$ can be defined by

$$2DFRFMT\{f(x, y, t, q)\} = F_{\alpha, \beta}(\xi, \eta, \lambda, \chi) = \langle f(x, y, t, q), K_{\alpha, \beta}(x, y, t, q, \xi, \eta, \lambda, \chi) \rangle \dots(4)$$

where, $K_{\alpha, \beta}(x, y, t, q, \xi, \eta, \lambda, \chi)$

$$= \sqrt{\frac{1 - icota}{2\pi}} e^{\frac{1}{2i\sin\alpha}[(x^2 + y^2 + \xi^2 + \eta^2)\cos\alpha - 2(x\xi + y\eta)]} e^{\frac{2i\pi i\lambda}{t\sin\theta} - 1} e^{\frac{2i\pi i\chi}{q\sin\theta} - 1} e^{\frac{\pi i}{\tan\theta}[\lambda^2 + \chi^2 + \log^2 t + \log^2 q]}$$

$$= C_{1\alpha} e^{iC_{2\alpha}[(x^2 + y^2 + \xi^2 + \eta^2)\cos\alpha - 2(x\xi + y\eta)]} t^{C_{1\theta}i\lambda - 1} q^{C_{1\theta}i\chi - 1} e^{C_{2\theta}i[\lambda^2 + \chi^2 + \log^2 t + \log^2 q]}$$

where $C_{1\alpha} = \sqrt{\frac{1 - icota}{2\pi}}$, $C_{2\alpha} = \frac{1}{2i\sin\alpha}$, $C_{1\theta} = \frac{2\pi}{\sin\theta}$,

$$C_{2\theta} = \frac{\pi}{\tan\theta}, 0 < \alpha < \frac{\pi}{2}, 0 < \theta < \frac{\pi}{2} \dots(5)$$

Right hand side of equation (4) has a meaning as the application of $f(x, y, t, q) \in E^*(\mathbb{R}^n)$ to

$$K_{\alpha, \beta}(x, y, t, q, \xi, \eta, \lambda, \chi) \in E.$$

It can be extended to the complex space as an entire function given by

$$2DFRFMT\{f(x, y, t, q)\} = F_{\alpha, \beta}(\xi', \eta', \lambda', \chi') = \langle f(x, y, t, q), K_{\alpha, \beta}(x, y, t, q, \xi', \eta', \lambda', \chi') \rangle \dots(6)$$

The right hand side is meaningful because for each $\xi', \eta', \lambda', \chi' \in \mathbb{C}^n$, $K_{\alpha, \beta}(x, y, t, q, \xi', \eta', \lambda', \chi') \in E$ as a function of x, y, t, q .

IV. APPLICATION OF TWO DIMENSIONAL FRACTIONAL FOURIER-MELLIN TRANSFORM

The kernel of two Dimensional Fractional Fourier-Mellin transform as

$$K_{\alpha, \beta}(x, y, t, q, \xi, \eta, \lambda, \chi) = C_{1\alpha} e^{iC_{2\alpha}[(x^2 + y^2 + \xi^2 + \eta^2)\cos\alpha - 2(x\xi + y\eta)]} t^{C_{1\theta}i\lambda - 1} q^{C_{1\theta}i\chi - 1} e^{C_{2\theta}i[\lambda^2 + \chi^2 + \log^2 t + \log^2 q]}$$

$$D_x D_y D_t D_q [K_{\alpha, \beta}(x, y, t, q, \xi, \eta, \lambda, \chi)] = C_{1\alpha} e^{iC_{2\alpha}(\xi^2 + \eta^2)\cos\alpha} D_x [e^{iC_{2\alpha}(x^2\cos\alpha - 2(x\xi))}] D_y [e^{iC_{2\alpha}(y^2\cos\alpha - 2(y\eta))}]$$

$$e^{C_{2\theta}i[\lambda^2 + \chi^2]} D_t [t^{C_{1\theta}i\lambda - 1} e^{C_{2\theta}i\log^2 t}] D_q [q^{C_{1\theta}i\chi - 1} e^{C_{2\theta}i\log^2 q}]$$

$$= \{C_{1\alpha} e^{iC_{2\alpha}(\xi^2 + \eta^2)\cos\alpha} e^{iC_{2\alpha}(x^2\cos\alpha - 2(x\xi))} iC_{2\alpha}[2x\cos\alpha - 2\xi] e^{iC_{2\alpha}(y^2\cos\alpha - 2(y\eta))} iC_{2\alpha}[2y\cos\alpha - 2\eta]\}$$

$$\{[e^{iC_{2\theta}[\lambda^2 + \chi^2]} t^{C_{1\theta}i\lambda - 1} e^{C_{2\theta}i\log^2 t} (iC_{2\theta}) \frac{2i\log t}{t} + e^{C_{2\theta}i\log^2 t} (C_{1\theta}i\lambda - 1)t^{C_{1\theta}i\lambda - 2}][q^{C_{1\theta}i\chi - 1} e^{C_{2\theta}i\log^2 q} (iC_{2\theta}) \frac{2i\log q}{q} + e^{C_{2\theta}i\log^2 q} (C_{1\theta}i\chi - 1)q^{C_{1\theta}i\chi - 2}]\}$$

$$= -4C_{2\alpha}^2(x\cos\alpha - \xi)(y\cos\alpha - \eta) C_{1\alpha} e^{iC_{2\alpha}(\xi^2 + \eta^2)\cos\alpha} e^{iC_{2\alpha}(x^2\cos\alpha + y^2\cos\alpha) - 2(x\xi + y\eta)} e^{iC_{2\theta}(\lambda^2 + \chi^2 + \log^2 t + \log^2 q)} t^{C_{1\theta}i\lambda - 1} q^{C_{1\theta}i\chi - 1} [(iC_{2\theta}) \frac{2i\log t}{t} + (C_{1\theta}i\lambda - 1)t^{-1}] [(iC_{2\theta}) \frac{2i\log q}{q} + (C_{1\theta}i\chi - 1)q^{-1}]$$

$$= -4C_{2\alpha}^2(x\cos\alpha - \xi)(y\cos\alpha - \eta) \frac{1}{tq} (2iC_{2\theta}\log t + C_{1\theta}i\lambda - 1)(2iC_{2\theta}\log q + C_{1\theta}i\chi - 1) C_{1\alpha} e^{iC_{2\alpha}[(x^2 + y^2 + \xi^2 + \eta^2)\cos\alpha - 2(x\xi + y\eta)]} t^{C_{1\theta}i\lambda - 1} q^{C_{1\theta}i\chi - 1} e^{iC_{2\theta}(\lambda^2 + \chi^2 + \log^2 t + \log^2 q)}$$

$$= -4 \frac{1}{4\sin^2\alpha} (x\cos\alpha - \xi)(y\cos\alpha - \eta) \frac{1}{tq} 2i(C_{2\theta}\log t + C_{1\theta}i\lambda - \frac{i}{2}) 2i(C_{2\theta}\log q + C_{1\theta}i\chi - \frac{i}{2}) K_{\alpha, \beta}(x, y, t, q, \xi, \eta, \lambda, \chi)$$

$$D_x D_y D_t D_q [K_{\alpha, \beta}(x, y, t, q, \xi, \eta, \lambda, \chi)] = (4\operatorname{cosec}^2\alpha)(x\cos\alpha - \xi)(y\cos\alpha - \eta) \frac{1}{tq} (C_{2\theta}\log t + C_{1\theta}i\lambda - \frac{i}{2}) K_{\alpha, \beta}(x, y, t, q, \xi, \eta, \lambda, \chi)$$

$$(C_{2\theta}\log q + C_{1\theta}i\chi - \frac{i}{2}) K_{\alpha, \beta}(x, y, t, q, \xi, \eta, \lambda, \chi) x^{-1}y^{-1}tq D_x D_y D_t D_q [K_{\alpha, \beta}(x, y, t, q, \xi, \eta, \lambda, \chi)] = (4\operatorname{cosec}^2\alpha)xy(\cos\alpha - \frac{\xi}{x})(\cos\alpha - \frac{\eta}{y}) \frac{1}{tq} (tC_{2\theta}\log t + tC_{1\theta}i\lambda - \frac{it}{2}) (qC_{2\theta}\log q + qC_{1\theta}i\chi - \frac{iq}{2}) K_{\alpha, \beta}(x, y, t, q, \xi, \eta, \lambda, \chi)$$

$$\Lambda_{x,y,t,q} K_{\alpha, \beta}(x, y, t, q, \xi, \eta, \lambda, \chi) = x^{-1}y^{-1}tq D_x D_y D_t D_q [K_{\alpha, \beta}(x, y, t, q, \xi, \eta, \lambda, \chi)] - (4\operatorname{cosec}^2\alpha)xy(\cos\alpha - \frac{\xi}{x})(\cos\alpha - \frac{\eta}{y}) \frac{1}{tq} (tC_{2\theta}\log t + tC_{1\theta}i\lambda - \frac{it}{2}) (qC_{2\theta}\log q + qC_{1\theta}i\chi - \frac{iq}{2}) K_{\alpha, \beta}(x, y, t, q, \xi, \eta, \lambda, \chi)$$

$$\Lambda_{x,y,t,q} = x^{-1}y^{-1}tq D_x D_y D_t D_q - (4\operatorname{cosec}^2\alpha)xy(\cos\alpha - \frac{\xi}{x})(\cos\alpha - \frac{\eta}{y}) \frac{1}{tq} (tC_{2\theta}\log t + tC_{1\theta}i\lambda - \frac{it}{2}) (qC_{2\theta}\log q + qC_{1\theta}i\chi - \frac{iq}{2})$$

$$\Lambda_{x,y,t,q} K_{\alpha, \beta}(x, y, t, q, \xi, \eta, \lambda, \chi) = (4\operatorname{cosec}^2\alpha) K_{\alpha, \beta}(x, y, t, q, \xi, \eta, \lambda, \chi)$$

$$\Lambda_{x,y,t,q} = (4\operatorname{cosec}^2\alpha)$$

$$\Lambda_{x,y,t,q} K_{\alpha, \beta}(x, y, t, q, \xi, \eta, \lambda, \chi) = (4\operatorname{cosec}^2\alpha) K_{\alpha, \beta}(x, y, t, q, \xi, \eta, \lambda, \chi)$$

$$\Lambda^2_{x,y,t,q} K_{\alpha, \beta}(x, y, t, q, \xi, \eta, \lambda, \chi) = (4\operatorname{cosec}^2\alpha)^2 K_{\alpha, \beta}(x, y, t, q, \xi, \eta, \lambda, \chi)$$

$$\Lambda^3_{x,y,t,q} K_{\alpha, \beta}(x, y, t, q, \xi, \eta, \lambda, \chi) = (4\operatorname{cosec}^2\alpha)^3 K_{\alpha, \beta}(x, y, t, q, \xi, \eta, \lambda, \chi)$$

$$\therefore \Lambda^k_{x,y,t,q} K_{\alpha,\beta}(x, y, t, q, \xi, \eta, \lambda, \chi) = (4 \operatorname{cosec}^2 \alpha)^k K_{\alpha,\beta}(x, y, t, q, \xi, \eta, \lambda, \chi)$$

Since the operator

$$\therefore \Lambda^k_{x,y,t,q} K_{\alpha,\beta}(x, y, t, q, \xi, \eta, \lambda, \chi) = (4 \operatorname{cosec}^2 \alpha)^k K_{\alpha,\beta}(x, y, t, q, \xi, \eta, \lambda, \chi)$$

Is obvious linear and continuous for $\alpha > 0, \theta > 0$

We have

$$\begin{aligned} & 2DFRFMT\{\Lambda^k_{x,y,t,q} f(x, y, t, q)\} \\ &= \langle \Lambda^k_{x,y,t,q} f(x, y, t, q), K_{\alpha,\beta}(x, y, t, q, \xi, \eta, \lambda, \chi) \rangle \\ &= \langle f(x, y, t, q), \Lambda^k_{x,y,t,q} K_{\alpha,\beta}(x, y, t, q, \xi, \eta, \lambda, \chi) \rangle \\ &= \langle f(x, y, t, q), (4 \operatorname{cosec}^2 \alpha)^k K_{\alpha,\beta}(x, y, t, q, \xi, \eta, \lambda, \chi) \rangle \end{aligned}$$

For all $f \in E'$ for $0 < \alpha < \frac{\pi}{2}, 0 < \theta < \frac{\pi}{2}$.

Adjoint Operator $\Lambda^*_{x,y,t,q}$

We define an operator $\Lambda^*_{x,y,t,q}: E' \rightarrow E$ using the relation

$$\begin{aligned} & \langle \Lambda^*_{x,y,t,q} [f(x, y, t, q)], \phi(x, y, t, q) \rangle \\ &= \langle [f(x, y, t, q)], \Lambda_{x,y,t,q} [\phi(x, y, t, q)] \rangle \end{aligned}$$

For all $f \in E'$ and $\phi \in E$. The operator $\Lambda^*_{x,y,t,q}$ is called the

adjoint operator of $\Lambda_{x,y,t,q}$ for each

$$k = 1, 2, 3, 4 \dots$$

We easily get,

$$\begin{aligned} & \langle (\Lambda^*_{x,y,t,q})^k f(x, y, t, q), K_{\alpha,\beta}(x, y, t, q, \xi, \eta, \lambda, \chi) \rangle \\ &= \langle f(x, y, t, q), (\Lambda^*_{x,y,t,q})^k K_{\alpha,\beta}(x, y, t, q, \xi, \eta, \lambda, \chi) \rangle \\ &= \langle f(x, y, t, q), (4 \operatorname{cosec}^2 \alpha)^k K_{\alpha,\beta}(x, y, t, q, \xi, \eta, \lambda, \chi) \rangle \\ &= (4 \operatorname{cosec}^2 \alpha)^k \langle f(x, y, t, q), K_{\alpha,\beta}(x, y, t, q, \xi, \eta, \lambda, \chi) \rangle \\ & 2DFRFMT\{(\Lambda^*_{x,y,t,q})^k f(x, y, t, q)\} = (4 \operatorname{cosec}^2 \alpha)^k \\ & \{2DFRFMT [f(x, y, t, q)]\} \\ & (\xi, \eta, \lambda, \chi) \dots (4.1) \end{aligned}$$

V. AN APPLICATION OF THE TWO DIMENSIONAL FRACTIONAL FOURIER-MELLIN TRANSFORM TO

Differential Equations

Solution of $P(\Lambda^*_{x,y,t,q})v = g$

$$\dots (5.1)$$

Consider, the differential equation $P(\Lambda^*_{x,y,t,q})v = g$

Where, $g \in E'$ and P is any polynomial of degree m .

Suppose that the equation (5.1) possesses a solution of v .

Applying $2DFRFMT$ to equation (5.1)

We have

$$2DFRFMT\{[P(\Lambda^*_{x,y,t,q})]v\} = 2DFRFMT\{g\}$$

$$\text{if } 2DFRFMT\{g\} = g^\wedge$$

$$\dots (5.2)$$

\therefore we can write

$$P(4 \operatorname{cosec}^2 \alpha) 2DFRFMT(v) = 2DFRFMT(g)$$

$$P(4 \operatorname{cosec}^2 \alpha) v^\wedge = g^\wedge \quad \text{by using (5.2)}$$

$$\dots (5.3)$$

If we further assume that the polynomial P is such that for $\epsilon > 0$,

$$\therefore |P(4 \operatorname{cosec}^2 \alpha)| < \epsilon < 0, \quad 0 < \alpha < \frac{\pi}{2}$$

Then under this assumption (5.3) gives

$$v^\wedge = [P(4 \operatorname{cosec}^2 \alpha)]^{-1} g^\wedge$$

$$\dots (5.4)$$

$$2DFRFMT(v) = g^\wedge [P(4 \operatorname{cosec}^2 \alpha)]^{-1}$$

Applying inversion of two dimensional fractional Fourier-Mellin transform to above equation we have

$$v = [2DFRFMT]^{-1} \left[\frac{g^\wedge}{P(4 \operatorname{cosec}^2 \alpha)} \right]$$

Hence proved.

VI. SOLUTION OF DIFFERENTIAL EQUATION FOR TWO DIMENSIONAL FRACTIONAL FOURIER-MELLIN TRANSFORM

Consider the differential equation

$$P(D)v = f(x, y, t, q) \dots (6.1)$$

Where $f \in E'$ and $P(D) = \sum_{|\beta| \leq m} a_\beta D^\beta$

is a linear differential operator of order m with constant coefficients.

Suppose that the equation (6.1) possesses a solution v .

Applying the two dimensional fractional Fourier-Mellin transform to (6.1) and using

$$\begin{aligned} & D_x^m D_t^n K_{\alpha,\beta}(x, y, t, q, \xi, \eta, \lambda, \chi) \\ &= \sum_{r=0}^m \sum_{s=0}^n C_{m,n} C_\alpha [x \operatorname{cosec} \alpha - \xi]^{m-2r} [y \operatorname{cosec} \alpha - \eta]^{n-2s} \\ & e^{(m-r)u + (n-s)v} \end{aligned}$$

$$\sum_{k=0}^l \sum_{h=0}^j ||j! \left(\frac{2\pi i}{\tan \theta} \right)^{l-k+j-h} C_k(t) C_k(q) (\log t)^{l-k} (\log q)^{h-j} e^{(l-k)u + (j-h)v}$$

$$\left(\frac{1}{t} \right)^{2l-k} \left(\frac{1}{q} \right)^{2h-j} P(\lambda) P(\chi) t^{\frac{2\pi i l}{\sin \theta} - 1} q^{\frac{2\pi i j}{\sin \theta} - 1}$$

where, $C_{m,n} = \frac{m!}{r!(m-2r)!} \frac{n!}{s!(n-2s)!} (2i)^{m-r+n-s}$

$$C_\alpha = \operatorname{cosec} \alpha^{r+s} (C_{2\alpha})^{m-r+n-s}$$

$$u = i C_{2\alpha} \{[(x)^2 + \xi^2] \operatorname{cosec} \alpha - 2(x)\xi\}$$

$$v = i C_{2\alpha} \{[(y)^2 + \eta^2] \operatorname{cosec} \alpha - 2(y)\eta\}$$

$$C_k(t) = \frac{l!}{(l-2k)!} \left[\frac{1 - \log t}{2 \log t} \right]^k$$

$$C_k(q) = \frac{j!}{(j-2h)!} \left[\frac{1 - \log q}{2 \log q} \right]^h$$

We have, $2DFRFMT [P(D)v] = 2DFRFMT [f] = f^\wedge$

(say)

We can reform them to the two dimensional fractional Fourier-Mellin transform and hence we get

$$P(x, y, t, q) v^\wedge = f^\wedge$$

Where $P(x, y, t, q)$ is polynomial in x, y, t, q ,

$$v^\wedge = 2DFRFMT [v]$$

Under the assumption that the polynomial P is such that

$$P\{D_x^m D_t^n K_{\alpha,\beta}(x, y, t, q, \xi, \eta, \lambda, \chi)\} > l > 0 \text{ for}$$

$$l_1, l_2, l_3, l_4 \dots l_n \in \mathbb{R}^n$$

$$v^\wedge = P\{D_x^m D_t^n K_{\alpha,\beta}(x, y, t, q, \xi, \eta, \lambda, \chi)\} f^\wedge$$

Applying inversion of two dimensional fractional

Fourier-Mellin transform to above equation we have

$$v = [2DFRFMT]^{-1} \left[\frac{f^\wedge}{P\{D_x^m D_t^n K_{\alpha,\beta}(x, y, t, q, \xi, \eta, \lambda, \chi)\}} \right]$$

VII. CONCLUSION

We have obtained the differential operator Λ and Λ^* . Using it we have solved the differential equation of the type $P(\Lambda^*_{x,y,t,q})v = g$. Also an application of two-dimensional fractional Fourier-Mellin transform to differential equation is presented.

REFERENCES

- [1] Stephane Derrode, Mohamed Daoudi Faouzi Ghorbel, "Invariant content-based image retrieval using a complete set of Fourier-Mellin descriptors",
- [2] Jun Wang, Zhiyong Xu, and Jianlin Zhang, "Image registration with hyperspectral data based on fourier-mellin transform", International Journal of Signal Processing Systems Vol. 1, No. 1 June 2013.

- [3] Tim Kazik , Ali Haydar Goktogan , “Visual odometry based on the Fourier-Mellin transform for a rover using a monocular ground-facing camera”,
- [4] Meisen Pan, Jianjun Jiang, Fen Zhang, Qiusheng Rong, “Medical Image Registration using Fourier-Mellin Transform”, Journal of Computational Information Systems 11: 3 (2015) 1141–1148.
- [5] V. D. Sharma, P. B. Deshmukh, “Inversion Theorem of two dimensional Fractional Fourier-Mellin Transform”, International Journal of Applied Computational Science & Mathematics, Vol-4, No.-1, 2014, pp-17-24.
- [6] V. D. Sharma, P. B. Deshmukh, “Adjoint Operators of Two Dimensional Fractional Fourier-Mellin Transform”, International Journal on Recent and Innovation Trends in Computing and Communication , Volume: 3 Issue: 12, pp 6514 – 6517.
- [7] V. D. Sharma, P. B. Deshmukh, “Analyticity of Two Dimensional Fractional Fourier-Mellin Transform”, International Journal of Engineering Sciences & Research Technology,5(3) March, 2016, pp 507-511.
- [8] V. D. Sharma, P. B. Deshmukh, “Convolution theorem for Two Dimensional Fractional Fourier-Mellin Transform”, International Journal of Pure & Applied Research in Engineering & Technology, 2016, Volume 4 (9), 54-63.
- [9] V. D. Sharma, P. D. Dolas, “Inversion Formula For Two Dimensional Generalized Fourier-Mellin Transform And It’s Application”, International Journal of Advanced Scientific and Technical Research December 2011, ISSUE 1 VOL 2.
- [10] R.Milanese, ,Cherbuliez, M.,Pun, T.: “Invariant Content-Based Image Retrieval Using the Fourier-Mellin Transform”. International conference on Advances in pattern Recognition Proceedings of ICAPR 98.

Dr. V. D. Sharma is currently working as an Assistant professor in the department of Mathematics, Arts, Commerce and Science College, Kiran Nagar, Amravati-444606 (M.S.) India. She has got 20 years of teaching and research experience. She has obtained her Ph.D. degree in 2007 from SGB Amravati University Amravati. Her field of interest is Integral Transforms. Six research students are working under her supervision. She has published more than 100 research articles.

P. B. Deshmukh is an Assistant professor in the department of Mathematics, IBSS College of Engineering, Amravati 444602 (M.S.) India. She has obtained her master degree in 2006 from Sant Gadage Baba Amravati University, Amravati. She has got 9 years of teaching experience. She has 15 research articles in journals to her credit.