

3-Quinolinecarbaldehyde thiosemicarbazones: Synthesis from N-arylacetamide, Characterization, Antibacterial and Antioxidant activities

Sreenivasulu Enumula, Satish Mudam, Khursheed Ahmed

Abstract— Four different Schiff bases of 3-quinolinecarbaldehydes are synthesized from differently substituted thiosemicarbazides. The structures of compounds are established using modern analytical technique FT-IR, ¹H-NMR and mass spectral data. Antimicrobial screening results are also presented against clinical isolates of pathogenic strains of *Staphylococcus aureus*, and *E. coli*. Simultaneously, antioxidant activities also have been carried out using DPPH radical scavenging properties to understand other functional applications of compounds. Depending on the structures of four synthesized compound, the compound **d** showed higher free radical neutralizing abilities as well as antimicrobial activities that may be due to the presence of 6-fluoro group of the quinoline which might be facilitating the flow of electrons through polarization of electron density of the quinoline ring.

Index Terms—3-quinolinecarbaldehyde, Thiosemicarbazide, Schiff's bases, antimicrobial, DPPH scavenging.

I. INTRODUCTION

In recent years nitrogen containing heterocyclic compounds like quinoline has received considerable attention due to their biological and pharmaceutical activities. Quinoline, also known as benzpyridine is a heterocyclic aromatic organic compound which has few anti-microbial properties but its derivatives are known to possess a considerable amount of effect on microbial strains[1-3].

Quinoline and its derivatives have always attracted the interests of both synthetic and biological chemists alike because of its diverse chemical and pharmacological properties. It is found in several natural compounds and pharmacologically active substances displaying a broad range of biological activities[3-5].

Literature survey had revealed that the chemistry of hydrazine derivatives such as thiosemicarbazide and its hydrazones is of immense interest owing to their wide synthetic and analytical applications and biological activities[6]. Thiosemicarbazides and their condensation products with different aldehydes display interesting biological activities and have therefore attracted considerable pharmaceutical interest. They have been evaluated for last 50 years as anticancer[7], antiHIV[8], antibacterial[9], antiviral[10] and antifungal due to their ability to diffuse

through the semipermeable membrane of cell lines[11-14]. They also play an important role in the regulation of plant growth[15]. Due to their abundance in plants and ease of synthesis, this class of compounds has generated great interest for possible therapeutic uses.

The important role displayed by quinoline and its derivatives for various therapeutic and biological activities prompted us to synthesize a new range of thiosemicarbazones. These are important synthetic conjugates that can be prepared with greater flexibility in reaction conditions as well as pharmacophores of varied nature. Such strategies can be used to create newer hybrid compounds with improved activity[16-17].

In the present report, following four poly substituted quinolinecarbaldehyde such as 2-hydroxy quinolinecarbaldehyde(**1**), 2-chloroquinolinecarbaldehyde (**2**), 6-fluoro-2-hydroxy-quinolinecarbaldehyde (**3**) and 6-fluoro-2-chloro-quinolinecarbaldehyde(**4**) are used with thiosemicarbazide for the syntheses of four (**a**, **b**, **c** and **d**) thiosemicarbazone.

II. EXPERIMENTAL

Solvents used for syntheses (acetone, methanol, ethanol and dichloromethane) were of AR grade and further dried before their use by standard procedures. The starting materials such as aniline, DMF, POCl₃, trifluoro acetic acid, acetic anhydride were obtained from SD-FCL Chemical Limited, Mumbai, India. N-aryl-acetamide, phosphoryl chloride and thiosemicarbazide were purchased from Sigma-Aldrich. All compounds were routinely checked by TLC on silica gel plates using petroleum ether/ethyl acetate as solvent system and the developed plates were visualized by UV light and iodine vapours. The detailed synthesis has been shown in Scheme-1.

Four different 3-quinoline carbaldehyde thiosemicarbazones **a**, **b**, **c** and **d** were synthesized through multistep reactions. Starting from cyclization of N-arylacetamide in the presence of Vilsmeier-Haack (DMF/POCl₃) reagent gives rise to 2-chloro-3-quinolinecarbaldehyde. Subsequent treatment by concentrated acid and preceded condensation with various acid hydrazides produce proposed hydrazones as reported by Sreenivasulu Enumula et al[18] (**Scheme – I**). Further structural characterization antimicrobial and antioxidant activities were reported in this work.

Sreenivasulu Enumula, Dept. of Chemistry & Post Graduate Centre, Pooa College of Arts, Science & Commerce, Camp, Pune – 411001, Affiliated to University of Pune, Pune, INDIA

Satish Mudam, Albany Molecular Research Inc., Hyderabad – 500078, INDIA

Khursheed Ahmed, Dept. of Chemistry & Post Graduate Centre, Abeda Inamdar Sr. College of Arts, Science & Commerce, Camp, Pune – 411001, Affiliated to University of Pune, Pune, INDIA and Poona College of Arts, Commerce and Science, Camp, Pune - 411001, Affiliated to University of Pune, Pune, INDIA

3-Quinolinecarbaldehyde thiosemicarbazones: Synthesis from N-arylacетamide, Characterization, Antibacterial and Antioxidant activities




Figure 1: Stepwise synthesis of thiosemicarbazones

The final products were purified on a silica gel chromatographic column using petroleum ether/ethyl acetate in proportion of 7:3 by V/V as solvent system. The purity of compounds was routinely checked by TLC on silica gel plates visualizing under UV light and iodine vapours. Melting points of the synthesized compounds were determined with open capillary tube on a VEEGO melting point apparatus. The ¹H-NMR was recorded on a SHIMADZU 600 MHz at Savitribai Phule Pune University, Pune while IR spectra were recorded on JASCO FT-IR-4100 spectrometer.

The chemically synthesized four compounds (**a**, **b**, **c** and **d**) were tested for antimicrobial activity. Strains of both Gram positive and Gram negative bacteria were used for experimentation. Amoxicillin (25 µg/mL⁻¹) was used as standard which shows a zone of inhibition of 8mm. The compounds were serially diluted and different dilutions were tested against *Staphylococcus aureus*, and *E. coli*.

The earlier studies attracted us to check antioxidant activities of these thiosemicarbazones. In the present work, we have used DPPH radical scavenging activities as per the procedure reported [19]. 2, 2-Diphenyl-1, 2 - picrylhydrazil (DPPH) is a stable free radical which has an unpaired valence electron at one atom of nitrogen bridge and scavenging of (DPPH) radical is the basis of popular DPPH antioxidant assay²⁰. The IC₅₀ values for even the standard antioxidants like ascorbic acid and butyrate hydroxytoluene (BHT) were also considered for the limiting concentrations. Aliquots of eight sample concentrations (1, 5, 10, 20, 25, 50, 100, 150 ml) prepared in methanol were taken in eight different test tubes. Samples were accurately dissolved in methanol to make the required concentration by dilution techniques. DPPH was weighed & dissolved to make 0.004% of DPPH solution. To dissolve homogeneously magnetic stirrer was used. After making the desired sample concentrations, 5ml of 0.004 % DPPH solution was applied on each test tube by micropipette to make a total volume of 10 ml. The room temperature was recorded & solutions were kept for 30 min to complete the reaction. DPPH was applied into blank test tube at the same time where methanol was taken as blank. After 30 minutes the absorbance of each test tube were taken by single beam spectrophotometer. IC₅₀ were measured from the plot of % inhibition verses concentration.

III. RESULTS AND DISCUSSION

All the intermediates were confirmed by reported physical constants. The structures of the synthesized compounds (**a**, **b**, **c** and **d**) were established on the basis of

physical and spectral data. The proposed structures are shown in the following **Table 1**.

Antimicrobial Studies:

The chemically synthesized four compounds (**a**, **b**, **c** and **d**) were tested for antimicrobial activities. Strains of both Gram positive and Gram negative bacteria were used for experimentation. Amoxicillin (25 µg/mL⁻¹) was used as standard which shows a zone of inhibition of 8mm. The compounds were serially diluted and different dilutions were tested against *Staphylococcus aureus*, and *E. coli*.

Comparison of inhibition zone diameter in mm for four compounds at a concentration 50, 100, 150 and 200 µg/ml respectively against Gram negative bacteria *E. Coli* is shown in **Table 2**. All four compounds lack antibacterial activity against Gram positive bacteria *S. aureus*. The values indicate that the antibacterial activity of the compounds increases logarithmically with an increase in concentration. Although all four the compounds show a similar type of inhibition pattern, the compound **d** exhibits more antibacterial activity as compared to others. Since the compounds show activity against *E. coli* as a representative organism from the Gram negative *Enterobacteriaceae* family. All four compounds can be screened for other enteric organisms who are known pathogens like *Salmonell*, *Pseudomonas*, *Vibrio* etc.

Table 2: Comparison of antimicrobial activities

No	Conc. (µg/ml)	a	b	c	d
1	50	8	8	8	10
2	100	10	9	10	11
3	150	12	9	12	13
4	200	13	12	14	15

Antioxidant activities:

The use of 2, 2-diphenyl-1-picryl-hydrazyl-hydrate (DPPH) assay provides an easy and rapid way to evaluate antioxidants by spectrophotometry²⁰. DPPH free radical method is an antioxidant assay based on electron-transfer process that produces a violet solution in ethanol. This free radical, stable at room temperature, is reduced in the presence of an antioxidant molecule (probably the synthesized conjugates) giving rise to colourless ethanol solution. From the values given in the following **Table 3**, it is seen that **c** and **d** have lowest IC₅₀ values compared to **a** and **b**. Based on such observations we can conclude that **c** and **d** are more potent antioxidant compounds than **a** and **b**.

One-electron oxidation of the compound **d** in 200 µg/ml solution may produce a phenoxyl radical at 2-hydroxy position of quinoline ring, indicating that the phenolic moiety of compound participates in free radical scavenging reactions. Depending on the structure, the compound **d** possesses higher free radical neutralizing abilities as well as antimicrobial activities that may be due to the presence of 6-fluoro group of the quinoline. This might facilitate the polarization of electron density of the molecule.

Table 3: DPPH radical scavenging activities in terms of IC₅₀ values of compounds

Compounds	IC ₅₀
a	2.4
b	4.0
c	5.6
d	6.4

Table 1: Spectral Characterization of synthesized compounds

SR. No.	Name Code	YIELD (%)	M. P. (°C)	Spectral Assignments
1	a	78%	202	FTIR (cm⁻¹): 710-770(C-Cl), 1647.56 (-C=N), 3195.86 (-NH), 2958.27 (-CH), 1660.41 (-C=S), 1057.03 (-N-N), 1500 to 1600 (Aromatic region). ¹H-NMR (dmsO) (δ, ppm): 9.77 ppm, 6.29 mm(s,1H,HC=N), 8.67 ppm, 6.72 mm (s, 1H, NH), 8.34 ppm, 6.64 mm (s, 1H, NH), 8.45ppm, 5.81 mm (d, 1h), 7.94-7.50, 6.12 + 6.33 + 6.32 + 6.31 mm (m, 4H), 3.84ppm, 13.00 mm (q, 2H), 1.38 ppm, 20.42 mm (t, 3H)
2	b	80%	196	FTIR (cm⁻¹): 1095(C-O), 1647.56 (-C=N), 3265.86 (-NH), 2958.27 (-CH), 1660.41 (-C=S), 1057.03 (-N-N), 3051.27 (-OH), 1500 to 1600 (Aromatic region), 2958.27 (-CH). ¹H-NMR (dmsO) (δ, ppm): 11.87 ppm, 2.03 mm (s, 1H, HC=N), 9.22 ppm, 2.07 mm (s, 1H-NH), 8.45ppm,2.12 mm (d, 1H), 8.52 ppm, 2.07 mm (s, 1H, -NH), 8.1-7.6 ppm, (4.17+2.17+2.32) mm (m, 4H),3.15 ppm, 6.79 mm (s, 3H)
3	c	70%	320	FTIR (cm⁻¹): 710-770 (C-Cl), 1647.56 (-C=N), 3265.86 (-NH), 2958.27 (-CH), 1660.41 (-C=S), 1057.03 (-N-N), 1500 to 1600 (Aromatic region). ¹H-NMR (CDCl₃) (δ, ppm): 12.02 ppm, 1.91 mm (s, 1HH, -OH), 11.70 ppm, 1.98 mm (s, 1H, HC=N), 8.68 ppm, 4.58 mm (s, 1H, -NH), 8.60 ppm, 4.58 mm (d, 1H), 8.28 ppm, 2.00 mm (s,1H, -NH), 7.6-7.2 ppm, (2.32+2.09+4.41) mm (m, 4H),3.05 ppm, 7.50 mm (s, 3H)
4	d	70%	202	FTIR (cm⁻¹): 710-770(C-Cl), 1647.56 (-C=N), 3195.86 (-NH), 2958.27 (-CH), 1660.41 (-C=S), 1057.03 (-N-N), 1500 to 1600 (Aromatic region). ¹H-NMR (dmsO) (δ, ppm): 11.91 ppm, 1.48 mm (s, 1H, HC=N), 9.17 pmm, 1.51 mm (s, 1H, -NH), 8.73 ppm, 1.65 mm (d, 1H), 8.50 ppm, 1.56 mm (s, 1H, -NH), 8.0-7.3 ppm, (1.64+3.67+0.75) mm (m, 4H), 3.15ppm, 5.36mm (s, 3H)

IV. CONCLUSION

In the present study, we have made an attempt to elaborate 3-Quinolinecarbaldehyde thiosemicarbazides as main pharmacophore. All the intermediates were confirmed by reported physical constants. The synthesized thiosemicarbazides are characterized by IR and ¹H-NMR. They are further tested for their in vitro antibacterial and antioxidant activities. All the synthesized compounds are moderately active when compared with standard Amoxicillin. Since the compounds show activity against E coli as a representative organism from the Gram negative Enterobacteriaceae family. All samples can be screened for other enteric organisms which are known pathogens like Salmonella, Pseudomonas, Vibrio etc. The results of antioxidant activities in terms of DPPH radical scavenging properties indicate the probable application of these compounds as antioncogenic agents in future research. The contest of development of new molecules has become more interesting and challenging with the advent in the field of drug design and discovery due to various multi resistant strains of pathogenic micro-organism. It has been observed that in this competition, pathogens have the upper hand since they can more easily evolve and develop a mechanism to fight the drug action. So, it is necessary to develop drugs which can easily act against the resistant strains.

ACKNOWLEDGMENT

The authors are grateful to the principal, Abeda Inamdar Senior College, Pune for extending the laboratory facilities. The authors extend their thanks to the Head, Department of microbiology for providing facilities to perform antibacterial assay.

REFERENCES

- [1] Aliasghar J., Khalili D., Clercq E.D., Salmi C. and Brunel J.M., Synthesis, Antibacterial, Antifungal and Antiviral Activity Evaluation of Some New bis-Schiff Bases of Isatin and Their Derivatives, *Molecules*, 2007, 12, 1720-1730.
- [2] Sari N., Arslan S., Logoglu E. and Sakiyan I., Antibacterial activities of some Amino acid Schiff bases, *Journal of Animal Science*, 2003, 16(2), 283-288.
- [3] Saidul I.M., Akhter F.M. and Bodruddoza M.A.K., Antineoplastic Activity of 2-oxo Benzylidene(3-oxo-Aniline)Cu(II) Ethylenediamine, *Pak. J. Biol Sci*, 2002, 5, 335-337.
- [4] Iana V., Emanuela T., Francesca M., and Fabio S., Aromatic Schiff bases and 2,3- disubstituted-1,3-thiazolidin-4-one derivatives as antiinflammatory agents, *Arkivoc*, 2004, 5, 364-74.
- [5] Matsubara H., Obara Y. and Kuba N., Biological activity of heterocyclic compounds II Blindness in baby chicks caused by administration of guanamines, guanides, and their related compounds, *Nipp. Nog. Kag. Kaishi.*, 1975, 49, 505-11.
- [6] Sperry J.B. and Wright D.L., Furans, thiophenes and related heterocycles in drug discovery, *Curr. Opini. Drug Disco. Develop*, 2005, 8, 723-40.
- [7] Su W., Zhang W., Wang T., Yan X. and Cai, J.D., Naphthane compound, derivative of Andrographolide and its medicinal application, *Fami. Zhu. Shen. Gong. Shu*, 2008, 33.
- [8] Winn M., Arendsen D., Dodge P., Dren A., Dunnigan D., Hallas R., Hwang K., Kyncl J., Lee Y., Plotnikoff N., Young P. and Zaugg H., Drugs derived from cannabinoids-5-delta6a,10a-Tetrahydrocannabinol and heterocyclic analogs containing aromatic side chains, *J. med. Chem.*, 1976, 19(4), 461-71.
- [9] Renault J., Giorgi S., Baron M., Mailliet P., Paoletti C., Cros S. and Voisin E., Heterocyclic Quinones. 4. A new highly cytotoxic drug: 6,7-bis(1-aziridinyl)-5,8-quinazolidinedione, *J. med. Chem.*, 1983, 26(12), 1715-1719.
- [10] Supuran C., Scozzafava A. and Andrea J., Carbonic Anhydrase Inhibitors. Part 541: Metal Complexes of Heterocyclic Sulfonamides: A New Class of Antiglaucoma Agents, *Metal-Based Drugs*, 1997, 4(6), 307-315.
- [1] Vicini P., Geronikaki A. and Incerti M., Aminoderivatives of benzothiazole/benzisothiazole heterocycles as intermediates in the synthesis of novel therapeutic agents, *Trudy Mezhdunarodnoi Konferentsii "Khimiya i Bio. Aktivnost Kislorod-i Serusoderz hashchikh Geterotsiklov"*, 2nd, Moscow, 2003, 1, 211-216.
- [2] Salvador J., Pinto R. and Silvestre S., Recent advances of pharmaceutical interest, *Current Org. Syn.*, 2009, 6(4), 426-470.

3-Quinolinecarbaldehyde thiosemicarbazones: Synthesis from N-arylacetamide, Characterization, Antibacterial and Antioxidant activities

- [3] Holan G., Mode of action of DDT- new aryl-alicyclic and heterocyclic insecticides, *Environ. Qua. Safety*, 1975, 3, 359-64.
- [4] Johnston M., Lohr J., Moes J., Solomon K. and Zaborski E., Toxicity of synergized and unsynergized nitromethylene heterocycle insecticide (SD 35651) to susceptible and resistant strains of *Musca domestica*, *J. eco. entomology*, 1986, 79(6), 1439-42.
- [5] Stehrer P. and Wolf H., Genotoxic evaluation of three heterocyclic Nmethylcarbamate pesticides using the mouse bone marrow micronucleus assay and the *Saccharomyces cerevisiae* strains D7 and D61.M., *Mutation res.*, 1995, 345(3-4), 111- 125.
- [6] Obara Y., Umemoto Y., and Matsubara H., Studies on the biological activity of heterocyclic compounds. Part IV. On the blinding activities and acute oral toxicities of aminobenzenes, pyrimidines, purines, amino-s-triazines and their related compounds in baby chicks, *Nipp. Nog. Kag. Kaishi.*, 1981, 55, 1205-1212.
- [7] Saczewski F. and Saczewski J., Applications of 2-chloro-4, 5-dihydroimidazole in heterocyclic and medicinal chemistry, *Trend. Hetero. Chem.*, 2003, 9, 19-31.
- [8] Sreenivasulu Enamula, Nusrat Shaikh, and Khursheed Ahmed, In Vitro Antibacterial Study of Synthesized 3-Quinolinecarbaldehyde through Vilsmeier Haack Cyclisation of N-Arylacetamide and their Hydrazones, *International Journal for Pharmaceutical Research Scholars*, 2014, 3(3), 26-32.
- [9] Afzal M., Al-oriquat G., Al-Hassan J.M., Muhammad N., Flavone glycosides from *Lawsonia inermis*. *Heterocycles*, 1980, 14, 1973-1976.
- [10] Dekkers J., Kooijman H., Kroon J., and Grech E., 2-Hydroxy-1,4-naphthoquinone, *Acta Crystallogr., Sect. C: Cryst. Struct. Commun.*, 1996, 52, 2896.